

**Math 280 Final Review Study Guide -
Problem 1 -**

1. Consider three position vectors (tails are the origin):

$$\vec{u} = \langle 1, 0, 0 \rangle$$

$$\vec{v} = \langle 4, 0, 2 \rangle$$

$$\vec{w} = \langle 0, 1, 1 \rangle$$

- (a) Find an equation of the plane passing through the tips of \vec{u} , \vec{v} , and \vec{w} .
- (b) Find an equation of the line perpendicular to the plane from part (a) and passing through the origin.

Solution:

- (a) Since the tails of the given vectors are at the origin, the tips of the vectors are the points $U = (1, 0, 0)$, $V = (4, 0, 2)$, and $W = (0, 1, 1)$, respectively. The plane containing the tips has $\vec{n} = \overrightarrow{UV} \times \overrightarrow{UW}$ as a normal vector. Since $\overrightarrow{UV} = \langle 3, 0, 2 \rangle$ and $\overrightarrow{UW} = \langle -1, 1, 1 \rangle$, the normal vector is

$$\vec{n} = \overrightarrow{UV} \times \overrightarrow{UW} = \langle -2, -5, 3 \rangle$$

Using $U = (1, 0, 0)$ as a point on the plane, an equation for the plane is

$$-2(x - 1) - 5(y - 0) + 3(z - 0) = 0$$

- (b) The line perpendicular to the plane in part (a) is parallel to the plane's normal vector. Thus, since $\langle -2, -5, 3 \rangle$ is parallel to the line and the origin $(0, 0, 0)$ is on the line, the vector equation for the line is

$$\vec{r}(t) = \langle 0, 0, 0 \rangle + t \langle -2, -5, 3 \rangle$$

Math 280 Final Review Study Guide
Problem 2

2. Consider the curve $\vec{\mathbf{r}}(t) = \langle t, t^3 \rangle$, $-\infty < t < \infty$.

- (a) Find the curvature $\kappa(t)$.
- (b) Find all values of t where $\kappa(t) = 0$.
- (c) Compute the limits

$$\lim_{t \rightarrow \infty} \kappa(t), \quad \lim_{t \rightarrow -\infty} \kappa(t)$$

(d) What do the limits in part (c) say about the curve $\vec{\mathbf{r}}(t)$?

Solution:

(a) By definition, the curvature of a curve parametrized by $\vec{\mathbf{r}}(t)$ is given by the formula

$$\kappa(t) = \frac{|\vec{\mathbf{r}}'(t) \times \vec{\mathbf{r}}''(t)|}{|\vec{\mathbf{r}}'(t)|^3}$$

The first two derivatives of $\vec{\mathbf{r}}(t)$ are $\vec{\mathbf{r}}'(t) = \langle 1, 3t^2 \rangle$ and $\vec{\mathbf{r}}''(t) = \langle 0, 6t \rangle$ and their cross product is $\vec{\mathbf{r}}'(t) \times \vec{\mathbf{r}}''(t) = 6t \hat{\mathbf{k}}$. Thus, the curvature of $\vec{\mathbf{r}}(t)$ is

$$\begin{aligned} \kappa(t) &= \frac{|\vec{\mathbf{r}}'(t) \times \vec{\mathbf{r}}''(t)|}{|\vec{\mathbf{r}}'(t)|^3}, \\ \kappa(t) &= \frac{6|t|}{\|\langle 1, 3t^2 \rangle\|^3}, \\ \kappa(t) &= \frac{6|t|}{(1 + 9t^4)^{3/2}} \end{aligned}$$

- (b) The curvature is 0 when $t = 0$.
- (c) The limits of $\kappa(t)$ as $t \rightarrow \pm\infty$ are

$$\lim_{t \rightarrow \pm\infty} \kappa(t) = \lim_{t \rightarrow \pm\infty} \frac{6|t|}{(1 + 9t^4)^{3/2}} = 0$$

(d) Lines are curves of zero curvature. Thus, the limits in part (c) suggest that $\vec{\mathbf{r}}(t)$ behaves linearly as $t \rightarrow \pm\infty$.


Math 280 Final Review Study Guide -
Problem 3 -

3. Given the function of two variables $G(x, y) = \sqrt{y - x^2}$

- (a) Determine the domain of G .
- (b) Sketch the level curves $G = 0$, $G = 1$, and $G = 2$ all on one coordinate grid. What kind of curves are they?
- (c) At the point $(1, 2)$, find the direction in which G has its maximum rate of increase. Also determine this maximum rate.

Solution:

- (a) The domain of G is the set of all pairs (x, y) such that $y - x^2 \geq 0$.



- (c) The direction of maximum rate of increase of $G(x, y)$ at the point $(1, 2)$ is, by definition,

$$\hat{\mathbf{u}} = \frac{\vec{\nabla} G(1, 2)}{\|\vec{\nabla} G(1, 2)\|}$$

The gradient of G is

$$\vec{\nabla} G = \langle G_x, G_y \rangle = \left\langle -\frac{x}{\sqrt{y-x^2}}, \frac{1}{2\sqrt{y-x^2}} \right\rangle$$

The value of $\vec{\nabla} G$ at the point $(1, 2)$ is $\vec{\nabla} G(1, 2) = \left\langle -1, \frac{1}{2} \right\rangle$ and its magnitude is $\|\vec{\nabla} G(1, 2)\| = \frac{\sqrt{5}}{2}$. Thus, the direction of maximum rate of increase of G at $(1, 2)$ is

$$\hat{\mathbf{u}} = \frac{\langle -1, \frac{1}{2} \rangle}{\frac{\sqrt{5}}{2}}$$

The maximum rate of increase, by definition, is $\vec{\nabla} G(1, 2) = \frac{\sqrt{5}}{2}$.

Math 280 Final Review Study Guide -
Problem 4 -

4. Find absolute maximum and minimum of the function $f(x, y) = xy - x$ over the region $R = \{x^2 + y^2 \leq 4\}$. Also, find the points where these extremes occur.

Solution: First, the region R is closed and bounded (i.e. compact) and f is defined at every point in R . Therefore, we are guaranteed to find absolute extrema. Next, we look for all critical points of f in R . These will be points for which the first derivatives of f vanish. Thus, we must solve the system of equations:

$$\begin{aligned}f_x &= y - 1 = 0, \\f_y &= x = 0\end{aligned}$$

which has $x = 0$ and $y = 1$ as the only solution. We must now determine the extreme values of f on the boundary of R which is the circle $x^2 + y^2 = 4$. We will resort to using the method of Lagrange multipliers to find these values. The following system of equations must then be solved:

$$\begin{aligned}f_x &= \lambda g_x, \\f_y &= \lambda g_y, \\g(x, y) &= 0\end{aligned}$$

where $g(x, y) = x^2 + y^2 - 4$. Evaluate the partial derivatives we then have

$$y - 1 = \lambda(2x), \tag{1}$$

$$x = \lambda(2y), \tag{2}$$

$$x^2 + y^2 = 4. \tag{3}$$

Dividing Equation (1) by Equation (2) and simplifying gives us

$$\begin{aligned}\frac{y - 1}{x} &= \frac{\lambda(2x)}{\lambda(2y)}, \\ \frac{y - 1}{x} &= \frac{x}{y}, \\ y(y - 1) &= x^2, \\ x^2 &= y^2 - y\end{aligned}$$

Substituting $y^2 - y$ for x^2 in Equation (3) and solving for x we get

$$\begin{aligned}x^2 + y^2 &= 4, \\y^2 - y + y^2 &= 4, \\2y^2 - y - 4 &= 0\end{aligned}$$

which has the two solutions

$$y_{1,2} = \frac{1 \pm \sqrt{33}}{4}$$

Let y_1 be the positive solution and y_2 the negative one. If $y = y_1$ then the corresponding x -values are $x_{11,12} = \pm\sqrt{y_1^2 - y_1}$. Similarly, if $y = y_2$ then the corresponding x -values are $x_{21,22} = \pm\sqrt{y_2^2 - y_2}$.

We must now evaluate $f(x, y)$ at the critical point $(0, 1)$ and at all critical points on the boundary of R .

$$\begin{aligned}f(0, 1) &= 0, \\f(x_{11}, y_1) &= x_{11}(y_1 - 1) = (y_1 - 1) \sqrt{y_1^2 - y_1} = \sqrt{y_1}(y_1 - 1)^{3/2} \\f(x_{12}, y_1) &= x_{12}(y_1 - 1) = -(y_1 - 1) \sqrt{y_1^2 - y_1} = -\sqrt{y_1}(y_1 - 1)^{3/2} \\f(x_{21}, y_2) &= x_{21}(y_2 - 1) = (y_2 - 1) \sqrt{y_2^2 - y_2} = \sqrt{-y_2}(1 - y_2)^{3/2} \\f(x_{22}, y_2) &= x_{22}(y_2 - 1) = -(y_2 - 1) \sqrt{y_2^2 - y_2} = -\sqrt{-y_2}(1 - y_2)^{3/2}\end{aligned}$$

A calculator would be useful here but isn't necessary. We can estimate $\sqrt{33}$ to be 5.75 using a linear approximation of $F(x) = \sqrt{x}$ about $x = 36$ giving us $y_1 \approx 1.6875$ and $y_2 \approx -1.1875$. One can then show that $f(x_{21}, y_2)$ is the absolute maximum and $f(x_{22}, y_2)$ is the absolute minimum of f on R .

Math 280 Final Review Study Guide
Problem 5


5. Consider the integral

$$\int_0^1 \int_{x^2}^1 x \cos y^2 \, dy \, dx.$$

- (a) Sketch the region of integration.
- (b) Reverse the order of integration properly.
- (c) Evaluate the integral from part (b).

Solution:

(a) The region of integration is sketched below.



(b) Upon switching the order of integration we obtain

$$\int_0^1 \int_0^{\sqrt{y}} x \cos y^2 \, dx \, dy$$

(c) Evaluating the above double integral we get

$$\begin{aligned} \int_0^1 \int_0^{\sqrt{y}} x \cos y^2 \, dx \, dy &= \int_0^1 \left[\frac{1}{2} x^2 \cos y^2 \right]_0^{\sqrt{y}} dy, \\ &= \frac{1}{2} \int_0^1 y \cos y^2 \, dy, \\ &= \frac{1}{2} \left[\frac{1}{2} \sin y^2 \right]_0^1, \\ &= \frac{1}{4} \sin(1) \end{aligned}$$

Math 280 Final Review Study Guide -
Problem 6 -

6. Consider the following vector field in space

$$\vec{\mathbf{F}} = \langle x + y, x + z, y \rangle.$$

- (a) Check that this field is conservative.
- (b) Find a potential of $\vec{\mathbf{F}}$.
- (c) Evaluate the following line integral

$$\int_C \vec{\mathbf{F}} \cdot d\vec{\mathbf{r}},$$

where C is a contour originating at $(0, 0, 0)$ and terminating at $(0, 1, 1)$.

Solution:

- (a) Let $P = x + y$, $Q = x + z$, and $R = y$. Given that $P_y = Q_x = 1$, $P_z = R_x = 0$, and $Q_z = R_y = 1$ we know that $\vec{\mathbf{F}}$ is conservative by the cross-partials test.
- (b) By inspection, a potential function for $\vec{\mathbf{F}}$ is $\varphi(x, y, z) = \frac{1}{2}x^2 + xy + yz$.
- (c) Using the Fundamental Theorem of Line Integrals, we obtain

$$\begin{aligned} \int_C \vec{\mathbf{F}} \cdot d\vec{\mathbf{r}} &= \varphi(0, 1, 1) - \varphi(0, 0, 0), \\ &= \left(\frac{1}{2}(0)^2 + 0 \cdot 1 + 1 \cdot 1 \right) - \left(\frac{1}{2}(0)^2 + 0 \cdot 0 + 0 \cdot 0 \right), \\ &= 1 \end{aligned}$$

Math 280 Final Review Study Guide -
Problem 7 -

7. Compute the circulation of the vector field

$$\vec{\mathbf{H}} = -y^3, x^3$$

over the boundary of the region $D = \{x^2 + y^2 \leq 1, y \geq 0\}$.

Solution: The boundary of D is a simple, closed curve oriented counter clockwise. Therefore, we may use Green's Theorem to compute the circulation:

$$\oint_{\partial D} \vec{\mathbf{H}} \bullet d\vec{\mathbf{r}} = \iint_D (Q_x - P_y) dA$$

Letting $P = -y^3$ and $Q = x^3$ we get $Q_x = 3x^2$ and $P_y = -3y^2$. Therefore, $Q_x - P_y = 3(x^2 + y^2)$. Since D is a half-disk, we will use polar coordinates to evaluate the double integral above. The integrand then becomes $3r^2$, $dA = r dr d\theta$, and the region D can be described as $\{0 \leq r \leq 1, 0 \leq \theta \leq \pi\}$. Thus, the circulation is

$$\begin{aligned} \oint_{\partial D} \vec{\mathbf{H}} \bullet d\vec{\mathbf{r}} &= \iint_D (Q_x - P_y) dA, \\ &= \int_0^\pi \int_0^1 3r^2 \cdot r dr d\theta, \\ &= \int_0^\pi \left[\frac{3}{4} r^4 \right]_0^1 d\theta, \\ &= \int_0^\pi \frac{3}{4} d\theta, \\ &= \frac{3\pi}{4} \end{aligned}$$

Math 280 Final Review Study Guide -
Problem 8 -

8. Compute the volume of the spherical wedge given in spherical coordinates by

$$W = \{ 1 \leq \rho \leq 2, 0 \leq \theta \leq \frac{\pi}{2}, 0 \leq \phi \leq \frac{\pi}{2} \}$$

Solution: Using spherical coordinates, the volume of the wedge is computed as follows

$$\begin{aligned} V &= \int_W 1 \, dV, \\ &= \int_0^{\pi/2} \int_0^{\pi/2} \int_1^2 \rho^2 \sin \phi \, d\rho \, d\phi \, d\theta, \\ &= \int_0^{\pi/2} \int_0^{\pi/2} \frac{1}{3} \rho^3 \sin \phi \Big|_1^2 \, d\phi \, d\theta, \\ &= \int_0^{\pi/2} \int_0^{\pi/2} \frac{7}{3} \sin \phi \, d\phi \, d\theta, \\ &= \int_0^{\pi/2} \left[-\frac{7}{3} \cos \phi \right]_0^{\pi/2} \, d\theta, \\ &= \int_0^{\pi/2} \frac{7}{3} \, d\theta, \\ &= \frac{7\pi}{6} \end{aligned}$$